



## Ibuprofen Dosage (Advil, Motrin)

Every 6 to 8 hours as needed for fever and pain

Weight (lbs)	Infants' Ibuprofen Concentrated Drops (50 mg)	Children's Ibuprofen Suspension Liquid (100 mg / 5 mL)	Junior Strength Tablet (100 mg)	Regular Strength Tablet (200 mg)
	<b>Dropper</b> (use dropper in mL)	<b>Teaspoon</b> (use dosing cup)	<b>Tablet</b>	<b>Tablet</b>
12 - 16	1.25 ml	2.5 ml (1/2 teaspoon)		
17 - 21	1.875 ml	3.75 ml (3/4 teaspoon)		
22 - 32	2.5 ml	5 ml (1 teaspoon)		
33 - 43	3.75 ml	7.5 ml (1 1/2 teaspoon)		
44 - 54	5ml	10 ml (2 teaspoons)	2 tablets	1 tablet
55 - 65		12.5 ml (2 1/2 teaspoons)	2 1/2 tablets	1 tablet
66 - 87		15 ml (3 teaspoons)	3 tablets	1 tablet
88+		20 ml (4 teaspoons)	4 tablets	2 tablets

**Ibuprofen is not recommended for patients under 6 months of age**  
**Do not exceed 4 doses of Ibuprofen in 24 hours**  
**If your child is receiving Ibuprofen for fever for longer than 24 hours, please call our office**