



Tylenol Dosage Chart

Every 4 to 6 hours as needed for fever and pain

Weight (lbs)	Infants Non-Concentrated Drops (160 mg / 5 ml)	Children's Suspension Liquid (160 mg / teaspoon)	Junior Strength Chewable Tablets (160 mg each)	Regular Strength Tablets (325 mg each)	Extra Strength Tablets (500 mg each)
	Syringe	Syringe or Teaspoon	Tablet	Tablet	Tablet
6 - 8	1.25 ml	1.25 ml (1/4 teaspoon)			
9 - 11	2 ml	2 ml			
12 - 17	2.5 ml	2.5 ml (1/2 teaspoon)			
18 - 23	3.75 ml	3.75 ml (3/4 teaspoon)			
24 - 29	5 ml (1 teaspoon)	5 ml (1 teaspoon)	1 tablet		
30 - 34	6.25				
35 - 46	7.5 (1 1/2 teaspoon)	7.5ml (1 1/2 teaspoon)	1 1/2 tablets		
47 - 74		10 ml (2 teaspoons)	2 tablets	1 tablet	
75+		15 ml (3 teaspoons)	3 tablets		1 tablet
<p>Do not administer Tylenol to children less than 6 weeks old</p> <p>Children under 4 months old need to be seen by your provider</p> <p>If your child is receiving Tylenol for fever for longer than 24 hours please call our office</p>					